

Europium anomalies constrain the mass of recycled lower continental crust

Ming Tang*, Roberta L. Rudnick, William F. McDonough, Richard M. Gaschnig, and Yu Huang

Department of Geology, University of Maryland, College Park, Maryland 20742, USA

ABSTRACT

Statistical analyses of Sm-Eu-Gd concentrations in more than 3000 samples from the upper, middle, and lower continental crust reveal that the enrichment of Eu in the lower continental crust cannot compensate for the Eu deficit in the upper and middle continental crust, leaving the bulk continental crust with a significant negative Eu anomaly. Because the building blocks of the continental crust (mantle-derived basalts or tonalitic slab melts) do not possess a negative Eu anomaly, removal of Eu-enriched lower continental crust is required. Using Sm-Eu-Gd systematics and a mass conservation model, at least $2.9^{+1.4}_{-0.5}$ (95% confidence) crustal masses ($\sim 6 \times 10^{22}$ kg) appear to have been lost to the mantle over Earth's history via lower crustal recycling. Such a lower crustal component in the mantle may reappear in some ocean island basalts that have positive Eu anomalies and unradiogenic Pb isotopes.

INTRODUCTION

The compositional differences within the continental crust are manifestations of intracrustal differentiation (Taylor and McLennan, 2008), which may occur in mature continental crust or precursor arc crust. It is widely accepted that the composition of the bulk continental crust is andesitic and is too evolved to be derived directly from the mantle (Arculus, 1981; Green et al., 2014; Rudnick, 1995). To explain the evolved crust composition, recycling of mafic to ultramafic lower crust has been proposed to occur through processes such as density foundering (Arndt and Goldstein, 1989; Kay and Mahlburg-Kay, 1991; Rudnick, 1995; Jull and Kelemen, 2001; Davidson and Arculus, 2005), subduction erosion (Clift et al., 2009; Lee, 2014, and references therein), or crustal subduction followed by reamination (Hacker et al., 2011).

Lower crustal recycling is poorly documented due to its non-observable nature in most circumstances (Taylor and McLennan, 2008). The geochemical evidence provided here supports the hypothesis that intracrustal differentiation followed by lower crustal recycling is important in crustal evolution. Although lower crustal recycling is mostly studied in Phanerozoic arc systems (e.g., Greene et al., 2006; Jagoutz and Schmidt, 2013), it may also occur in cratons (e.g., Gao et al., 2004). Our geochemical approach using Sm-Eu-Gd systematics constrains the total mass of recycled lower continental crust (LCC) throughout Earth history, independent of the tectonic environments, processes, and timing of recycling.

Sm-Eu-Gd SYSTEMATICS IN THE CONTINENTAL CRUST

Europium is present in two valence states (+2 and +3). During intracrustal differentiation, Eu is fractionated from Sm and Gd, as Eu^{2+} substitutes for Ca^{2+} and Na^{+} in feldspar (Ren, 2004), which is stable under crustal pressure-temperature (P - T) conditions. The magnitude of Eu enrichment or depletion in a sample is calculated as a "Eu anomaly" relative to the neighboring rare earth elements Sm and Gd:

$$\frac{\text{Eu}}{\text{Eu}^*} = \frac{\text{Eu}_N}{\sqrt{(\text{Sm}_N)(\text{Gd}_N)}}, \quad (1)$$

where the subscript N indicates normalization to chondritic values (Sun and McDonough, 1989). Feldspar left behind in the LCC during intracrustal differentiation causes a Eu deficit in the upper continental crust

(UCC) and enrichment in the LCC (Taylor and McLennan, 2008). However, whether the current deep crustal reservoir compensates the Eu deficit in the UCC is unclear. Among the published compositional models for the bulk continental crust (BCC) (plotted in Appendix 1 in the GSA Data Repository¹), all but one have negative Eu anomalies of varying magnitude (0.78–0.97), while only Taylor and McLennan (2008) proposed $\text{Eu}/\text{Eu}^* = 1$ for the BCC.

Here we evaluate Eu/Eu^* in the UCC, middle continental crust (MCC), and LCC using an updated data compilation following that of Huang et al. (2013). This data set has three subsets: high- P granulite facies rocks, amphibolite facies rocks, and shales + loess + tillites. For simplicity, we assume that the bulk compositions of samples in the three data sets represent those of the LCC, MCC, and UCC, respectively. We have applied no weighting of samples based on their bulk compositions nor have we used any geophysical information regarding the bulk composition of the deep crust. For our calculation, we used only samples that have Sm, Eu and Gd concentration data, comprising 970 samples from the LCC, 1702 samples from the MCC, and 411 samples from the UCC. The amphibolite and granulite facies data compilations include samples having both meta-sedimentary and meta-igneous protoliths. We excluded 10% of the samples with high and low Sm and Gd concentrations (removing 2.5% from the top and bottom of the Sm concentration distribution, then repeating the process for Gd), and then removed another 10% of samples with high and low Eu/Eu^* . This double filter, which eliminates 20% of the samples, helps to reduce the chance of incorporating atypical samples or bad analyses without shifting the results significantly (<10%).

Table 1 lists the mean Sm, Eu, and Gd concentrations in UCC, MCC, and LCC, the number of samples that went into the calculations after filtering, the values from Rudnick and Gao (2003), and the calculated mean concentrations of Sm, Eu, and Gd in the BCC, which are made using a bootstrap method with 100,000 times resampling of the compositional data set. This technique allows estimation of the bulk composition and associated uncertainties with all samples taken into account. The mass proportions of UCC, MCC, and LCC are based on global seismic data, as described by Huang et al. (2013), and the uncertainties were simulated using a Monte Carlo scheme (details in Appendix DR1). Uncertainties or biases associated with the sampling of the UCC, MCC, and LCC were not taken into account in these calculations. Although there are fewer UCC samples than MCC and LCC samples, sediments are powerful predictors of the bulk rare earth element (REE) composition of the UCC, as REEs are insoluble and are transported quantitatively from the site of weathering to the site of deposition (Taylor and McLennan, 2008).

Our results, along with previous estimates tabulated here, suggest that the average UCC and MCC have negative Eu anomalies, while the average LCC has a small positive Eu anomaly. To reconcile the Eu/Eu^* mass balance in the absence of lower crustal recycling, the Eu/Eu^* of the modern LCC would need to be at least ~ 1.7 (using our estimated Sm-Eu-Gd concentrations in the BCC and Sm-Gd in the LCC), and this seems unlikely in light of the existing data (Fig. 1). Although one could argue that the data compiled here may be biased toward lower Eu/Eu^* relative to the true com-

¹GSA Data Repository item 2015244, Appendix DR1 (method description and sensitivity tests) and Appendix DR2 (compiled data sets), is available online at www.geosociety.org/pubs/ft2015.htm, or on request from editing@geosociety.org or Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.

*E-mail: tangmyes@gmail.com

TABLE 1. MEAN CONCENTRATIONS OF Sm, Eu, AND Gd IN THE CONTINENTAL CRUST AND CANDIDATES OF RECYCLED LOWER CONTINENTAL CRUST

	UCC	MCC	LCC	BCC
Mass (%)*	35.9 ± 6.0	33.5 ± 4.3	30.6 ± 3.4	100
This work†				
n‡	328	1362	776	
Sm	5.3 ± 0.2	4.1 ± 0.1	3.6 ± 0.2	4.4 ± 0.2
Eu	1.08 ± 0.03	1.13 ± 0.03	1.22 ± 0.04	1.14 ± 0.03
Gd	4.5 ± 0.1	4.1 ± 0.1	3.7 ± 0.1	4.1 ± 0.1
Eu/Eu*	0.67 ± 0.01	0.84 ± 0.01	1.02 ± 0.02	0.82 ± 0.04
Rudnick and Gao (2003)				
Sm	4.7 ± 0.3	4.6	2.8	3.9
Eu	1 ± 0.1	1.4	1.1	1.1
Gd	4 ± 0.3	4	3.1	3.7
Eu/Eu*	0.72	0.96	1.14	0.89
Candidates for Recycled LCC				
	JS13	K14	RG03	This study**
Sm	1.04	1.03	2.8	2.7 ± 0.2
Eu	0.55	0.47	1.1	1.00 ± 0.06
Gd	1.77	1.09	3.1	2.9 ± 0.2
Eu/Eu*	1.24	1.36	1.14	1.10 ± 0.03
($m_{\text{recycled LCC}}/m_{\text{BCC}}$)	2.6	2.0	1.9	2.9 ± 1.1/–0.9

Note: The estimates of Rudnick and Gao (2003) are listed for comparison. The uncertainties in the estimates from this work are at the 95% confidence level. The uncertainties in the upper continental crust (UCC) values recommended by Rudnick and Gao (2003) are 1 σ standard deviation. MCC—middle continental crust; LCC—lower continental crust; BCC—bulk continental crust. $m_{\text{recycled LCC}}/m_{\text{BCC}}$ —mass proportion of recycled LCC relative to present-day continental crust. JS13—Arc delaminants from Jagoutz and Schmidt (2013). K14—Talkeetna arc lower crustal cumulates from Kelemen et al. (2014). RG03—LCC from Rudnick and Gao (2003).

*Mass proportions of the UCC, MCC, and LCC are from Huang et al. (2013).

†Concentrations given in ppm by weight.

‡After filtering for extreme values (see text).

**Means of granulite facies samples with Mg# > 50 and SiO₂ < 54%.

position of the LCC, we note that as Eu/Eu* increases in granulites, Eu concentration decreases (Fig. 2), so a more mafic (higher Mg#) lower crust with higher Eu/Eu* would have less leverage on the overall Eu/Eu* of the crust due to its lower REE concentration (see Appendix DR1 for sensitivity tests of assumed LCC composition). That is probably the reason that the Rudnick and Gao (2003) BCC, which incorporates a relatively mafic, high-Eu/Eu* LCC, still has a significant negative Eu anomaly (Table 1).

THE Eu DILEMMA

We assume, as many have (e.g., Taylor and McLennan, 2008; Gao et al., 1998), that the original crust produced by mantle melting has Eu/Eu* = 1. However, rather than possessing no Eu anomaly, the modern BCC shows significant Eu depletion relative to Sm and Gd (Table 1). These observations require that materials with positive Eu anomalies be transferred from the continental crust to the mantle. Our observations cannot constrain these processes, but can provide limits on the composition and mass of the recycled crust.

CALCULATING THE MASS OF THE RECYCLED LOWER CRUST

Generally, crustal recycling takes two forms: sediment subduction and LCC recycling (via gravitationally driven processes, e.g., density foundering, subduction erosion, relamination, etc.). Sediment subduction returns upper crustal materials to the mantle. Terrigenous sediments, which dominate the REE budget of global subducting sediments (Plank, 2014), have negative Eu anomalies, and thus cannot account for the Eu deficit in the continental crust. By contrast, recycling of the LCC is a plausible mechanism for decreasing the bulk crust Eu/Eu*, as the LCC is observed to be enriched in Eu.




Figure 1. Histogram showing the distribution of Eu/Eu* values in granulite facies samples (n = 970), compared with the Eu/Eu* of lower continental crust (LCC) required to make the bulk continental crust Eu/Eu* = 1 (black assuming the LCC has the Sm and Gd concentrations from this study). Kernel density is an estimate of probability function. Inset shows chondrite-normalized rare earth element patterns for the upper continental crust (UCC, solid gray curve), middle continental crust (MCC, dashed gray curve), and LCC (black curve) from Rudnick and Gao (2003).




Figure 2. Correlations between Eu/Eu*, Eu concentration and Mg# in granulite facies samples. The mean Eu/Eu*, Eu concentration and Mg# represent incremental averages. Each increment consists of 100 samples. High-Mg# samples have high Eu/Eu* but low Eu concentrations.

In our modeling, we make the following assumptions.

- (1) The present-day bulk continental crust has the composition provided in Table 1 and discussed herein.
- (2) Juvenile crust has a Eu/Eu* = 1.0.
- (3) The loss of upper crustal mass due to sediment subduction removes material with Eu/Eu* < 1.0 from the continental crust.
- (4) The Eu/Eu* of recycled lower crust can be characterized using the following four compositional models: (1) an estimate of recycled intra-oceanic arc lower crust (Jagoutz and Schmidt, 2013); (2) an estimate of high-density cumulates in arc lower crust (Kelemen et al., 2014); (3) an estimate of the recycled LCC from this work (see the following); and (4) the Rudnick and Gao (2003) estimate of the modern LCC.

The latter two of these assumptions bear further discussion. Given the complexities associated with sediment subduction (i.e., the degree to which sediments melt in the subarc environment and how this may influ-




Figure 3. Mass proportion of recycled lower continental crust (LCC; x-axis) needed to compensate the Eu depletion in the modern bulk continental crust (BCC) relative to an original crust with $\text{Eu}/\text{Eu}^* = 1$. By mass conservation, the modern BCC plus the recycled LCC equals the initial juvenile crust. The field of gray dots is generated by Monte Carlo resampling, and represents the predicted distribution using our estimate for the recycled LCC derived from granulite facies samples with $\text{Mg}\# > 50$ and $\text{SiO}_2 < 54\%$ (vertical black solid and dotted lines are the projected mean value and 95% confidence interval, respectively, of the mass proportion of recycled LCC). Curve JS13 represents the model outcome using Jagoutz and Schmidt's (2013) Kohistan arc estimate for the recycled lower crust; curve K14 is calculated using Talkeetna arc high-density lower crustal cumulates; curve RG03 uses the LCC from Rudnick and Gao (2003).

(1) Arc lower crustal delaminates. Jagoutz and Schmidt (2013) calculated the composition of missing negatively buoyant cumulates from the crustal section of the Kohistan arc using a major element mass balance approach.

(2) Arc lower crustal cumulates. Kelemen et al. (2014) calculated the average composition of Talkeetna arc high density lower crustal cumulates that may be subject to foundering.

(3) Mafic LCC granulites. We estimate the Sm-Eu-Gd composition of recycled lower crust based on the average compositions of granulites with $\text{Mg}\# > 50$ ($\text{Mg}\# = \text{molar ratio of Mg to Mg + Fe}$) and $\text{SiO}_2 < 54\%$ (double filter applied, as described here), which limits the recycled component to mafic compositions. The density of these mafic granulites, with an average SiO_2 and $\text{Mg}\#$ of 48% and 64, respectively, can increase by $\sim 10\%$ during eclogitization (Austheim, 1987) and exceed that of the underlying mantle (Krystopowicz and Currie, 2013). Although high-Mg# granulites, on average, have more positive Eu anomalies (Fig. 2), this is counterbalanced by the fact that high-Mg# granulites have low average Eu concentrations (Fig. 2).

(4) We also considered the LCC composition of Rudnick and Gao (2003) as an end member scenario.


The mass of recycled LCC is calculated using mass conservation equations for Sm-Eu-Gd (i.e., the Eu/Eu^* in modern BCC + recycled LCC = bulk juvenile crust) (Appendix DR1). Using the Sm-Eu-Gd compositions for the BCC estimated in this work, we calculated the mass of the recycled LCC to be at least $2.9^{+1.1}_{-0.9}$ (95% confidence) times the mass of the modern continental crust (Table 1; and Fig. 3). The results are not significantly different whether using the recycled lower crust compositions of arc cumulates (Jagoutz and Schmidt, 2013; Kelemen et al., 2014) or the Rudnick and Gao (2003) LCC (Fig. 3). Such consistency is not surprising given the anti-correlation between Eu/Eu^* and Eu concentration in various estimates of the recycled LCC (Table 1), similar to the observation in granulites (Fig. 2). This trade-off between Eu/Eu^* and Eu concentration makes the model relatively insensitive to the type of samples one chooses to calculate the composition of the recycled LCC. Our model is sensitive to the composition of BCC, which is subject to the uncertainties in sampling of deep crust. However, we show in Appendix DR1 that the BCC Eu/Eu^* is insensitive to the greatest unknown; i.e., the LCC composition.

The recycled LCC/BCC mass fraction required by our results over Earth history is within uncertainty of most previous estimates, all but one of which are based on regional case studies. Using Eu/Eu^* and Sr/Nd, Gao et al. (1998) estimated 1–2 crustal masses (37–82 km) of LCC were recycled in central east China. In the Mesozoic Sierra Nevada (California, USA) continental arc, 60%–75% of the original juvenile basalt (i.e., 1.5–3.0 crustal masses) is estimated to have been removed from the arc lower crust (Lee et al., 2007). This same ratio calculated by Jagoutz and Schmidt (2013) for the Kohistan arc ranges from 1.3 to 2.4 crustal masses.

ence their REE patterns [Plank, 2005], and whether they are relaminated to the lower crust [Behn et al., 2011]), we ignore sediment subduction in our mass balance model. Based on assumption 4, ignoring the subducted sediment flux leads to a minimum estimate of the mass of recycled lower crust.

The composition of the recycled LCC is difficult to assess except that these components may be denser than bulk crust, and possibly mantle peridotite (a density greater than peridotite is not a precondition for recycling, as mafic granulites may transform to high-density eclogite upon subduction or during crustal thickening), given their mafic to ultramafic bulk compositions. We therefore used four different lower crustal compositional estimates as candidates for the recycled lower crust (Table 1):

Figure 4. A: Distribution of Eu/Eu^* values in primitive ocean island basalts (OIBs) with $\text{MgO} > 10\%$ (wt%). **B:** $^{206}\text{Pb}/^{204}\text{Pb}$ versus Eu/Eu^* for OIB locality averages for OIBs with $\text{MgO} > 10\%$ (to avoid any influence from shallow magma differentiation); locality averages of Pb isotopes are calculated using all samples from each ocean island system. OIB data are from GeoRoc (georoc.mpch-mainz.gwdg.de/georoc). Average mid-oceanic ridge basalt (MORB) Eu/Eu^* is from our unpublished global primitive MORB data, and average MORB Pb isotopes are from Gale et al. (2013). LCC—lower continental crust.



A somewhat lower estimate of recycled lower crust (25%–60% by mass foundering of arc lower crust, which translates into > 0.3–1.5 modern crustal masses) was obtained by Plank (2005) using Th/La systematics of modern arcs and average continental crust composition. However, as she pointed out, this range is a minimum estimate.

The recycled LCC mass ($\sim 6 \times 10^{22}$ kg) may form a significant reservoir within the mantle. It has a positive Eu anomaly and low μ ($^{238}\text{U}/^{204}\text{Pb}$) (e.g., Kramers and Tolstikhin, 1997), unradiogenic Sr and Nd isotopes, and radiogenic Os isotopes (e.g., Lee, 2014). Average ocean island basalts (OIBs) also have a bulk positive Eu anomaly (Fig. 4A), which correlates negatively with $^{206}\text{Pb}/^{204}\text{Pb}$ (Fig. 4B). This negative correlation may reflect mixing between mantle components and recycled LCC with positive Eu anomalies and low $^{206}\text{Pb}/^{204}\text{Pb}$. Recycled LCC has been suggested to explain the geochemical characteristics of EM I type (enriched mantle I) OIB, typified by Kerguelen Plateau samples (e.g., Frey et al., 2002), which show pronounced positive Eu anomalies (mean $\text{Eu}/\text{Eu}^* = 1.12$) and unradiogenic Pb isotopes (mean $^{206}\text{Pb}/^{204}\text{Pb} = 18.3$). The recycled LCC, together with the modern continental crust, may balance the depleted mantle in the Sr and Nd isotope systems if the recycling occurred relatively early in Earth's history.

CONCLUDING REMARKS

Using available REE data for upper, middle, and lower crustal samples, we show that the bulk continental crust has a significant negative Eu anomaly ($\text{Eu}/\text{Eu}^* = 0.82 \pm 0.04$, 95% confidence), a result consistent with most previous estimates of Eu/Eu^* in the continental crust. Assuming that the original crust extracted from the mantle has no Eu anomaly, recycling of Eu-enriched LCC (mass proportion, $m_{\text{recycled LCC}} / m_{\text{BCC}}$ of $2.9^{+1.1}_{-0.9}$, 95% confidence) is required to account for the Eu deficit.

These findings affirm the important roles that intra-crustal differentiation and lower crustal recycling play in the formation and evolution of the continental crust. Moreover, they characterize the mass and composition of a component in the mantle (which could be spatially dispersed or concentrated) that plays a significant role in the chemical budget of the bulk silicate Earth. Consequently, models of growth of the continental crust (mass versus time) must consider that the total original crustal mass fraction could be a factor of four greater than the present mass of continental crust.

ACKNOWLEDGMENTS

This work was supported by National Science Foundation grants EAR-0948549, EAR-1067983 and EAR-1321954. Discussions with Michael N. Evans, Kang Chen, and Michael Brown, and constructive review comments from Peter Kelemen, Richard J. Arculus, Esteban Gazel, Jon Davidson, and an anonymous reviewer were very helpful in refining the ideas and model presented here.

REFERENCES CITED

- Arculus, R.J., 1981, Island arc magmatism in relation to the evolution of the crust and mantle: *Tectonophysics*, v. 75, p. 113–133, doi:10.1016/0040-1951(81)90212-2.
- Arndt, N.T., and Goldstein, S.L., 1989, An open boundary between lower continental crust and mantle: Its role in crust formation and crustal recycling: *Tectonophysics*, v. 161, p. 201–212, doi:10.1016/0040-1951(89)90154-6.
- Austrheim, H., 1987, Eclogitization of lower crustal granulites by fluid migration through shear zones: *Earth and Planetary Science Letters*, v. 81, p. 221–232, doi:10.1016/0012-821X(87)90158-0.
- Behn, M.D., Kelemen, P.B., Hirth, G., Hacker, B.R., and Massonne, H.-J., 2011, Diapirs as the source of the sediment signature in arc lavas: *Nature Geoscience*, v. 4, p. 641–646, doi:10.1038/ngeo1214.
- Clift, P.D., Schouten, H., and Vannucchi, P., 2009, Arc-continent collisions, sediment recycling and the maintenance of the continental crust, in Cawood, P.A., and Kröner, A., eds., *Earth Accretionary Systems in Space and Time: Geological Society of London Special Publication 318*, p. 75–103, doi:10.1144/SP318.3.
- Davidson, J.P., and Arculus, R.J., 2005, The significance of Phanerozoic arc magmatism in generating continental crust, in Brown, M., and Rushmer, T., eds., *Evolution and Differentiation of the Continental Crust: Cambridge, UK, Cambridge University Press*, p. 156–158.
- Frey, F.A., Weis, D., Borisova, A.Y., and Xu, G., 2002, Involvement of continental crust in the formation of the Cretaceous Kerguelen Plateau: New Perspectives from ODP Leg 120 Sites: *Journal of Petrology*, v. 43, p. 1207–1239, doi:10.1093/petrology/43.7.1207.

- Gale, A., Dalton, C.A., Langmuir, C.H., Su, Y., and Schilling, J.-G., 2013, The mean composition of ocean ridge basalts: *Geochemistry, Geophysics, Geosystems*, v. 14, p. 489–518, doi:10.1029/2012GC004334.
- Gao, S., Zhang, B.-R., Jin, Z.-M., Kern, H., Ting-Chuan, L., and Zhao, Z.-D., 1998, How mafic is the lower continental crust?: *Earth and Planetary Science Letters*, v. 161, p. 101–117, doi:10.1016/S0012-821X(98)00140-X.
- Gao, S., Rudnick, R.L., Yuan, H.L., Liu, X.M., Liu, Y.S., Xu, W.L., Ling, W.L., Ayers, J., Wang, X.C., and Wang, Q.H., 2004, Recycling lower continental crust in the North China craton: *Nature*, v. 432, p. 892–897, doi:10.1038/nature03162.
- Green, D.H., Hibberson, W.O., Rosenthal, A., Kovács, I., Yaxley, G.M., Falloon, T.J., and Brink, F., 2014, Experimental study of the influence of water on melting and phase assemblages in the upper mantle: *Journal of Petrology*, v. 55, p. 2067–2096, doi:10.1093/petrology/egu050.
- Greene, A.R., DeBari, S.M., Kelemen, P.B., Blusztajn, J., and Clift, P.D., 2006, A detailed geochemical study of island arc crust: The Talkeetna arc section, south-central Alaska: *Journal of Petrology*, v. 47, p. 1051–1093, doi:10.1093/petrology/egl002.
- Hacker, B.R., Kelemen, P.B., and Behn, M.D., 2011, Differentiation of the continental crust by reamination: *Earth and Planetary Science Letters*, v. 307, p. 501–516, doi:10.1016/j.epsl.2011.05.024.
- Huang, Y., Chubakov, V., Mantovani, F., Rudnick, R.L., and McDonough, W.F., 2013, A reference Earth model for the heat-producing elements and associated geoneutrino flux: *Geochemistry, Geophysics, Geosystems*, v. 14, p. 2003–2029, doi:10.1002/ggge.20129.
- Jagoutz, O., and Schmidt, M.W., 2013, The composition of the foundered complement to the continental crust and a re-evaluation of fluxes in arcs: *Earth and Planetary Science Letters*, v. 371–372, p. 177–190, doi:10.1016/j.epsl.2013.03.051.
- Jull, M., and Kelemen, P.B., 2001, On the conditions for lower crustal convective instability: *Journal of Geophysical Research*, v. 106, no. B4, p. 6423–6446, doi:10.1029/2000JB900357.
- Kay, R.W., and Mahlburg-Kay, S., 1991, Creation and destruction of lower continental crust: *Geologische Rundschau*, v. 80, p. 259–278, doi:10.1007/BF01829365.
- Kelemen, P.B., Hanghøj, K., and Greene, A.R., 2014, One view of the geochemistry of subduction-related magmatic arcs, with an emphasis on primitive andesite and lower crust, in Rudnick, R.L., ed., *The crust: Treatise on geochemistry volume 4 (second edition): Oxford, UK, Elsevier*, p. 749–806, doi:10.1016/B978-0-08-095975-7.00323-5.
- Kramers, J.D., and Tolstikhin, I.N., 1997, Two terrestrial lead isotope paradoxes, forward transport modelling, core formation and the history of the continental crust: *Chemical Geology*, v. 139, p. 75–110, doi:10.1016/S0009-2541(97)00027-2.
- Krystopowicz, N.J., and Currie, C.A., 2013, Crustal eclogitization and lithosphere delamination in orogens: *Earth and Planetary Science Letters*, v. 361, p. 195–207, doi:10.1016/j.epsl.2012.09.056.
- Lee, C.-T.A., 2014, Physics and chemistry of deep continental crust recycling, in Rudnick, R.L., ed., *The crust: Treatise on geochemistry volume 4 (second edition): Oxford, UK, Elsevier*, p. 423–456, doi:10.1016/B978-0-08-095975-7.00314-4.
- Lee, C.-T.A., Morton, D.M., Kistler, R.W., and Baird, A.K., 2007, Petrology and tectonics of Phanerozoic continent formation: From island arcs to accretion and continental arc magmatism: *Earth and Planetary Science Letters*, v. 263, p. 370–387, doi:10.1016/j.epsl.2007.09.025.
- Plank, T., 2005, Constraints from thorium/lanthanum on sediment recycling at subduction zones and the evolution of the continents: *Journal of Petrology*, v. 46, p. 921–944, doi:10.1093/petrology/egi005.
- Plank, T., 2014, The chemical composition of subducting sediments, in Rudnick, R.L., ed., *The crust: Treatise on geochemistry volume 4 (second edition): Oxford, UK, Elsevier*, p. 607–629, doi:10.1016/B978-0-08-095975-7.00319-3.
- Ren, M., 2004, Partitioning of Sr, Ba, Rb, Y, and LREE between alkali feldspar and peraluminous silicic magma: *American Mineralogist*, v. 89, p. 1290–1303.
- Rudnick, R.L., 1995, Making continental crust: *Nature*, v. 378, p. 571–578, doi:10.1038/378571a0.
- Rudnick, R., and Gao, S., 2003, Composition of the continental crust, in Rudnick, R.L., ed., *The crust: Treatise on geochemistry: Oxford, UK, Elsevier*, p. 1–64, doi:10.1016/B0-08-043751-6/03016-4.
- Sun, S.-S., and McDonough, W.F., 1989, Chemical and isotopic systematics of oceanic basalts: Implications for mantle composition and processes, in Saunders, A.D., and Norry, M.J., eds., *Magmatism in the Ocean Basins: Geological Society of London Special Publication 42*, p. 313–345, doi:10.1144/GSL.SP.1989.042.01.19.
- Taylor, S.R., and McLennan, S., 2008, *Planetary Crusts: Their Composition, Origin and Evolution: Cambridge, UK, Cambridge University Press*, 401 p.

Manuscript received 22 January 2015

Revised manuscript received 7 June 2015

Manuscript accepted 9 June 2015

Printed in USA

Geology

Europium anomalies constrain the mass of recycled lower continental crust

Ming Tang, Roberta L. Rudnick, William F. McDonough, Richard M. Gaschnig and Yu Huang

Geology 2015;43:703-706
doi: 10.1130/G36641.1

Email alerting services click www.gsapubs.org/cgi/alerts to receive free e-mail alerts when new articles cite this article

Subscribe click www.gsapubs.org/subscriptions/ to subscribe to *Geology*

Permission request click <http://www.geosociety.org/pubs/copyrt.htm#gsa> to contact GSA

Copyright not claimed on content prepared wholly by U.S. government employees within scope of their employment. Individual scientists are hereby granted permission, without fees or further requests to GSA, to use a single figure, a single table, and/or a brief paragraph of text in subsequent works and to make unlimited copies of items in GSA's journals for noncommercial use in classrooms to further education and science. This file may not be posted to any Web site, but authors may post the abstracts only of their articles on their own or their organization's Web site providing the posting includes a reference to the article's full citation. GSA provides this and other forums for the presentation of diverse opinions and positions by scientists worldwide, regardless of their race, citizenship, gender, religion, or political viewpoint. Opinions presented in this publication do not reflect official positions of the Society.

Notes